arXiv:1108.1220v1 [hep-ph] 4 Aug 2011

Tensorial NSI and Unparticle physics in neutrino-electron scattering

J. Barranco[*]
Instituto de Astronomía,
Universidad Nacional Autonoma de México,
México, DF 04510, Mexico

A. Bolaños[†]
Facultad de Ciencias Físico Matemáticas,
Benemérita Universidad Autónoma de Puebla,
Apartado Postal 1152, Puebla, Puebla, México.

E. A. Garcés[‡] and O. G. Miranda[§]
Departamento de Física, Centro de Investigación y de Estudios Avanzados del IPN,
Apdo. Postal 14-740 07000 México, D F, Mexico

T. I. Rashba[¶]
Max-Planck-Institute for Solar System Research,
Katlenburg-Lindau, 37191, Germany[**]

Abstract

We have analyzed the neutrino electron scattering off electrons in order to obtain constraints on tensorial couplings. We have studied the non standard interactions formalism as well as the unparticle stuff case. For our analysis we have focused on the recent TEXONO collaboration results. We have obtained new constraints to possible tensorial couplings in both new physics formalisms.

---

[*] Electronic address: barranco@astroscu.unam.mx

[†] Electronic address: azucena@fcfm.buap.mx

[‡] Electronic address: egarces@fis.cinvestav.mx

[§] Electronic address: Omar.Miranda@fis.cinvestav.mx

[¶] Electronic address: timur@mppmu.mpg.de

[**] On leave from IZMIRAN, Institute of Terrestrial Magnetism, Ionosphere and Radio Wave Propagation of

---

the Russian Academy of Sciences, 142190, Troitsk, Moscow region, Russia

I. INTRODUCTION

Searching for new physics has been highly motivated by the neutrino oscillation physics which implies non-zero neutrino mass [1]. In fact, neutrino oscillations are the first clue of physics beyond the standard model and has motivated a new generation of neutrino experiments as well as the development of phenomenological and theoretical research on models beyond the Standard Model [2, 3] . The future generation of neutrino experiments put the field as very promising in the search for new physics.

A very useful tool for the phenomenological study of these types of new physics is the formalism of non standard interactions (NSI). This formalism can parametrize a wide range of well motivated models of neutrino mass and, at the same time, gives constraints that are model independent. In the NSI formalism, operators with (V±A) Lorentz structure have been widely studied in the literature, giving stringent limits in neutrino-electron interactions when solar and reactor data is analyzed. However, tensorial non standard neutrino interactions have not been studied in detail in the neutrino sector.

A very different type of new physics that also can give rise to tensorial interactions is the recently proposed unparticle physics. In this extension, particles couple to a hidden conformal sector [4, 5] which could be probed in future experiments.

Unparticle physics can be directly produced in accelerator experiments. One way to probe their direct production is searching for signatures of missing energy in the detectors. There is another way to study their effects, which is through low energy processes mediated by the unparticle stuff.

The study of neutrino electron scattering has been of interest for elementary particle physics since their first measurement at the GARGAMELLE bubble chamber [6]. Recently, neutrino detectors using reactor neutrinos in a Short Baseline have taken profit of the intense and pure source of $\bar{\nu}_e$ to probe for physics beyond the Standard Model [7–9].

Neutrino electron scattering has been recently measured with improved accuracy by using reactor antineutrinos. The TEXONO collaboration has reported recent results [9, 10] and that has motivated us to study tensorial interactions in the framework of NSI as well as in the unparticle physics scheme. Here we introduce the results of such an analysis. We will see that it is possible to obtain strong constraints to the parameters of both approaches.

Our paper is organized as follows: In Section II we introduce the NSI formalism and present the corresponding neutrino electron cross Section, while in Section III we present the analog discussion for the unparticle case. The description of our analysis for the TEXONO case is shown in Section IV. Finally our conclusions are given in Section V.

II. TENSORIAL NSI IN NEUTRINO-ELECTRON SCATTERING

It is a common characteristic of many extensions of the Standard Model to introduce new interactions that can be parametrized with the help of an effective Lagrangian. The most studied case regards with $V - A$ extensions of the Standard Model. It is well known that in this case the effective four fermion Lagrangian takes the form: [11–15]

$$-\mathcal{L}^{eff}_{\rm V-A} = \varepsilon^{fP}_{\alpha\beta} 2\sqrt{2} G_F (\bar{\nu}_\alpha \gamma_\rho L \nu_\beta)(\bar{f}\gamma^\rho P f), \tag{1}$$

where f is a first generation SM fermion: e, u or d, and $P = L$ or R are the chiral projectors. There is plenty of literature studying either current constraints on the neutrino NSI with electrons [16–18] and quarks [19, 20] as well as future perspectives for long baseline neutrino experiments [21, 22] as well as in reactor low energy neutrino experiments [23].

In particular for the neutrino electron scattering which has the advantage of being a purely leptonic process and therefore is free from QCD uncertainties the neutrino electron cross section for the NSI case is given by

$$\begin{aligned}\frac{d\sigma(E_\nu, T)}{dT} &= \frac{2G_F^2 m_e}{\pi}[(\tilde{g}_L^2 + \sum_{\alpha\neq e} |\epsilon^{eL}_{\alpha e}|^2) + \\ &+ (\tilde{g}_R^2 + \sum_{\alpha\neq e} |\epsilon^{eR}_{\alpha e}|^2)\left(1 - \frac{T}{E_\nu}\right)^2 - (\tilde{g}_L\tilde{g}_R + \sum_{\alpha\neq e} |\epsilon^{eL}_{\alpha e}||\epsilon^{eR}_{\alpha e}|) m_e \frac{T}{E_\nu^2}],\end{aligned} \tag{2}$$

with $\tilde{g}_L = g_L + \epsilon^{eL}_{\alpha\alpha}$ and $\tilde{g}_R = g_R + \epsilon^{eR}_{\alpha\alpha}$. The standard model constants are given by $g_L = \frac{1}{2} + \sin^2\theta_W$, $g_R = \sin^2\theta_W$.

On the other hand, limited attention has been given to possible tensor interactions. This may be motivated by the $(V - A)$ structure of the Standard Model, however, with neutrino physics entering into a precision era it could be a good moment to study this kind of interactions with more detail.

Previous study to this interaction in neutrino electron scattering has been given in different articles [24] and there have been some constraints reported in the literature [25].



FIG. 1: Comparison between Standard-Model and tensor NSI differential cross section.

The Lagrangian describing this type of interaction is given by

$$-\mathcal{L}_{\mathrm{T}}^{eff}=\varepsilon_{\alpha\beta}^{fT}2\sqrt{2}G_F(\bar{\nu}_\alpha\sigma^{\mu\nu}\nu_\beta)(\bar{f}\sigma_{\mu\nu}f), \tag{3}$$

with $\sigma_{\mu\nu}=\gamma_\mu\gamma_\nu+\gamma_\nu\gamma_\mu$. In this case, for the tensorial contribution the amplitude for the process is given by

$$\mid M\mid^2= g_T^2\frac{G_F^2}{2}128m_e^2(4E_\nu^2+T^2-(4E_\nu+m_e)T) \tag{4}$$

and therefore the differential NSI neutrino electron cross section takes the form:

$$\frac{d\sigma_T^{NSI}}{dT}=\frac{\mid M\mid^2}{64\pi m_e E_\nu^2}=g_T^2\frac{4G_F^2m_e}{\pi}\Big[\Big(1-\frac{T}{2E_\nu}\Big)^2-\frac{m_eT}{4E_\nu^2}\Big]. \tag{5}$$

In this case, g_T parametrizes the strength of the tensorial NSI coupling. In order to have an idea about the tensorial NSI interaction, we plot the differential cross section versus the energy and compare it with the Standard Model prediction, Fig (1) .

III. NEUTRINO-ELECTRON SCATTERING MEDIATED BY TENSORIAL UNPARTICLE INTERACTIONS

At energies above certain Λ, a hidden sector operator $\mathcal{O}_{UV}$ of dimension d_{UV} could couple to the SM operators $\mathcal{O}_{SM}$ of dimension d_{SM} via the exchange of heavy particles of mass M

$$\mathcal{L}_{UV}=\frac{\mathcal{O}_{UV}\mathcal{O}_{SM}}{M^{d_{UV}+d_{SM}-4}}\,. \tag{6}$$

The hidden sector becomes scale invariant at Λ and then the interactions become of the form

$$\mathcal{L}_{\mathcal{U}} = C_{\mathcal{O}_{\mathcal{U}}} \frac{\Lambda^{d_{UV}-d}}{M^{d_{UV}+d_{SM}-4}} \mathcal{O}_{\mathcal{U}} \mathcal{O}_{SM} \,, \tag{7}$$

where $\mathcal{O}_{\mathcal{U}}$ is the unparticle operator of scaling dimension d in the low energy limit and $C_{\mathcal{O}_{\mathcal{U}}}$ is a dimensionless coupling constant. Therefore the unparticle sector can appear at low energies in the form of new massless fields coupled very weakly to the SM particles.

Scalar and vectorial unparticle interactions have been studied previously in the context of neutrino electron scattering [10, 26, 27]. In what follows we will concentrate on the case of tensorial interactions.

In the low energy regime, the effective interactions for the tensor unparticle operators have the form [28, 29]

$$\frac{-i}{4} \frac{\kappa_T}{\Lambda_{\mathcal{U}}^{d_{\mathcal{U}}}} \bar{f} \left(\gamma_\mu \overset{\leftrightarrow}{D}_\nu + \gamma_\nu \overset{\leftrightarrow}{D}_\mu \right) \psi_f \; \mathcal{O}_{\mathcal{U}}^{\mu\nu} \text{ and } \frac{\kappa_T}{\Lambda_{\mathcal{U}}^{d_{\mathcal{U}}}} \mathcal{F}_{\mu\alpha} \mathcal{F}_{\nu}^{\alpha} \, \mathcal{O}_{\mathcal{U}}{}^{\mu\nu}, \tag{8}$$

where $\mathcal{F}_{\mu\alpha}$ denotes the gauge field strength. From here we can find that the unparticle propagator for the tensorial field is given by

$$[\mathcal{A}_{\mathcal{F}}(P^2)]_{\mu\nu,\rho\sigma} = \frac{\mathcal{A}_d}{2\sin(d\pi)} (-P^2)^{d-2} T_{\mu\nu,\rho\sigma}(P), \tag{9}$$

where

$$\mathcal{A}_d = \frac{16\pi^{5/2}}{(2\pi)^{2d}} \frac{\Gamma(d+1/2)}{\Gamma(d-1)\Gamma(2d)} \tag{10}$$

and the tensor operator $T_{\mu\nu\rho\sigma}$ is given by

$$T_{\mu\nu\rho\sigma} = \frac{1}{2} \left\{ \pi_{\mu\rho}\pi_{\nu\sigma} + \pi_{\mu\sigma}\pi_{\nu\rho} - \frac{2}{3}\pi_{\mu\nu}\pi_{\rho\sigma} \right\}, \tag{11}$$

with

$$\pi_{\mu\nu} = -g_{\mu\nu} + \frac{P_\mu P_\nu}{P^2}. \tag{12}$$

If we consider the case of neutrino electron scattering we will arrive to the amplitude

$$M_T = \frac{f(d)}{\Lambda_U^{2d-2}} \left\{ \bar{\nu}(k')\sigma^{\mu\nu}\nu(k) \right\} \left(-(P)^2 - i\varepsilon \right)^{d_u - 2} T_{\mu\nu\rho\sigma} \left\{ e(p')\sigma^{\rho\sigma} e(p) \right\}, \tag{13}$$

where

$$f(d) = \frac{\lambda_{i\nu}^{\alpha\beta} \lambda_{if}}{2\sin(d\pi)} A_d. \tag{14}$$

In what follows we will use the definition for the couplings $\lambda = \sqrt{\lambda_{i\nu}^{\alpha\beta}\lambda_{if}}$ and we will fix the scale $\Lambda = 1$ TeV. From this amplitude, it is possible to obtain the differential cross section



FIG. 2: Comparison between Standard Model and tensor unparticle differential cross section for $\Lambda = 1$ TeV.

for neutrino electron scattering for the tensorial unparticle interactions. The computations of this cross section become more transparent if we decompose it into a electric part which takes the form:

$$\frac{d\sigma_T}{dT} = \frac{f(d)^2}{\pi\Lambda_U^{4d-4}} 2^{2d-3} m_e^{2d-3} T^{2d-4} \Big[\Big(1 - \frac{T}{2E_\nu}\Big)^2 - \frac{mT}{2E_\nu^2}\Big] \tag{15}$$

and a magnetic contribution that is given by

$$\frac{d\sigma_T}{dT} = \frac{f(d)^2}{\pi\Lambda_U^{4d-4}} 2^{2d-2} m_e^{2d-3} T^{2d-4} \Big(1 - \frac{T}{2E_\nu}\Big)^2. \tag{16}$$

In order to obtain the total cross section, this expressions should be added to the Standard Model prediction

$$\frac{d\sigma(\bar{\nu}_e)}{dT} = \frac{2G_F^2 m_e}{\pi}[g_R^2 + g_L^2(1 - \frac{T}{E_\nu})^2 - g_R g_L \frac{m_e T}{E_\nu^2}]. \tag{17}$$

We show in Fig. (2) the comparison between the Standard Model prediction for the differential cross section and the unparticle case, we can see that the expectations to obtain a good constraint from this process at low energies are encouraging.

IV. LIMITS ON TENSOR INTERACTIONS FROM THE TEXONO EXPERIMENT

Among the most recent reactor neutrino experiments, the TEXONO collaboration has published results on the cross section for the neutrino electron scattering [10, 30] using the Kuo Sheng 2.9 GW reactor as an antineutrino source that provides an average flux of $6.4 \times 10^{12} cm^{-2} s^{-1}$. Even though the collaboration has made use of three different detectors, we will focus on the CsI(Tl) detector data in order to obtain constraints for the tensor interactions both for the NSI and unparticle cases.

In order to obtain a constraint on the tensorial parameters we have computed the expected number of events for the TEXONO detector in the case of a NSI or unparticle interaction given by Eqs. (5) and (16,17) respectively and compute the integral

$$N_i = K \int_{T_i}^{T_{i+1}} \int_{E_\nu} \frac{d\sigma}{dT} \frac{d\phi(\bar{\nu}_e)}{dE_\nu} dE_\nu dT, \tag{18}$$

where the factor K accounts for the time exposure and the number of electron targets and $d\phi(\bar{\nu}_e)/dE_\nu$ is the neutrino spectrum which we have parametrized as the exponential of a polynomial order five as has been recently discussed in the literature [31]. We have also considered the relative abundances of each radioactive isotope in the nuclear reactor, ^{235}U(98%), ^{238}U(1.5%), and ^{239}Pu(0.4%). The electron recoil energy is divided into ten bins, T_i, running from 3 to 8 MeV.

Once we have computed the theoretical expected events per bin we can compute the χ^2 function

$$\chi^2 = \sum_{i=1} \left[\frac{N_{expt}(i) - [N_{NSI,U}(i)]}{\Delta_{stat}(i)} \right]^2 , \tag{19}$$

where $N_{NSI,U}(i)$, is the calculated event rate in the ith energy data bin for the Tensorial NSI or the unparticle cases, $N_{expt}(i)$ is the observed event rate for the corresponding energy bin, and $\Delta_{stat}(i)$ is the statistical uncertainty of the associated measurement.

The results of our analysis are shown in Figures (3) and (4). We can see that for the NSI case the constraint on the tensorial coupling gives the bound, at 90 % C. L., $g_T \leq 0.20$ which is much better than the previously reported constraint by the LAMPF collaboration [25]. We obtained the value of $\chi^2_{min} = 5.43$ for 9 d.o.f. We have verified explicitly that previous reactor experiments give weaker constraints than those obtained here using the TEXONO



FIG. 3: $\Delta\chi^2$ for the tensorial NSI coupling g_T in the TEXONO experiment.



FIG. 4: limits at 90 % CL for the tensorial unparticle parameters d and λ from our analysis of the recent TEXONO data.

data. For the unparticle case we have shown the 90 % CL region for the parameter λ and d. In this case we obtained $\chi^2_{min} = 5.16$ for 8 d.o.f. We have also extracted the constraints from the solar neutrino analysis reported in reference [32] (following analogous assumptions to those discussed in ref. [27]) and plotted the result in the same Figure (4) in order to show the interplay between different analysis; as can be seen the results from our analysis are more restrictive for values of $d > 2.03$.

experiment	g_T
TEXONO 90 % CL	0.20
LAMPF 90 % CL	0.379
Stellar energy loss	$0.06 - 3.6$

TABLE I: Limits on the tensorial coupling g_T from previous analysis and from the TEXONO experiment analyzed in this work.

V. DISCUSSION AND SUMMARY

Neutrino physics is entering into a precision era that could give important guidance about new physics beyond the Standard Model.

In this article we have concentrated in the case of tensorial couplings that could give a signal in reactor neutrino experiments. We have studied in particular the recent TEXONO results on antineutrino electron scattering. The tensorial interactions have been studied both in the framework of NSI and for the unparticle case. We have found new constraints that are stronger than previous laboratory constraints. These results can be summarized in Table I for the NSI case where we show the previous laboratory result from the LAMPF experiment [25]. Besides, we also show the astrophysical estimates that come from stellar energy loss [33, 34]. For the case of an unparticle tensor interaction, we have found that our constraints are more restrictive than previous analysis for values of $d > 2.03$.

As can be seen the results are encouraging and future neutrino electron scattering experiments could give even stronger constraints. Another possible place to search for this type of interaction in the future could be the coherent neutrino nuclei scattering that is also part of the TEXONO low energy neutrino physics program [35] and other proposals [36–38].

VI. ACKNOWLEDGMENTS

We would like to thank Muhammed Deniz and Celio Moura for useful discussions. This work has been supported by CONACyT.

---

[1] T. Schwetz, M. Tortola and J. W. F. Valle, New J. Phys. **13**, 063004 (2011) [arXiv:1103.0734 [hep-ph]].

[2] H. Nunokawa, S. J. Parke and J. W. F. Valle, Prog. Part. Nucl. Phys. **60**, 338 (2008) [arXiv:0710.0554 [hep-ph]].

[3] R. N. Mohapatra and A. Y. Smirnov, Ann. Rev. Nucl. Part. Sci. **56**, 569 (2006) [arXiv:hep-ph/0603118].

[4] H. Georgi, Phys. Rev. Lett. **98**, 221601 (2007) [arXiv:hep-ph/0703260].

[5] H. Georgi, Phys. Lett. B **650**, 275 (2007) [arXiv:0704.2457 [hep-ph]].

[6] F. J. Hasert *et al.* [Gargamelle Neutrino Collaboration], Nucl. Phys. **B73**, 1-22 (1974). F. J. Hasert *et al.* [Gargamelle Neutrino Collaboration], Phys. Lett. **B46**, 138-140 (1973).

[7] Z. Daraktchieva *et al.* [MUNU Collaboration], Phys. Lett. B **615**, 153 (2005) [arXiv:hep-ex/0502037].

[8] H. T. Wong *et al.* [TEXONO Collaboration], Phys. Rev. D **75**, 012001 (2007) [arXiv:hep-ex/0605006].

[9] M. Deniz *et al.* [TEXONO Collaboration], Phys. Rev. D **81**, 072001 (2010) [arXiv:0911.1597 [hep-ex]].

[10] M. Deniz *et al.* [TEXONO Collaboration], Phys. Rev. D **82**, 033004 (2010) [arXiv:1006.1947 [hep-ph]].

[11] V. D. Barger, R. J. N. Phillips and K. Whisnant, Phys. Rev. D **44**, 1629 (1991).

[12] S. Bergmann, M. M. Guzzo, P. C. de Holanda, P. I. Krastev and H. Nunokawa, Phys. Rev. D **62**, 073001 (2000) [arXiv:hep-ph/0004049].

[13] P. Huber and J. W. F. Valle, Phys. Lett. B **523**, 151 (2001) [arXiv:hep-ph/0108193].

[14] Z. Berezhiani and A. Rossi, Phys. Lett. B **535** 207 (2002), Z. Berezhiani, R.S. Raghavan and A. Rossi, Nucl. Phys. B **638** 62 (2002)

[15] S.Davidson, C. Pena-Garay, N. Rius and A. Santamaria JHEP **303**, 11 (2003);

[16] J. Barranco, O. G. Miranda, C. A. Moura and J. W. F. Valle, Phys. Rev. D **73**, 113001 (2006) [arXiv:hep-ph/0512195].

[17] J. Barranco, O. G. Miranda, C. A. Moura and J. W. F. Valle, Phys. Rev. D **77**, 093014 (2008) [arXiv:0711.0698 [hep-ph]].

[18] A. Bolanos, O. G. Miranda, A. Palazzo, M. A. Tortola and J. W. F. Valle, Phys. Rev. D **79**, 113012 (2009) [arXiv:0812.4417 [hep-ph]].

[19] F. J. Escrihuela, M. Tortola, J. W. F. Valle and O. G. Miranda, Phys. Rev. D **83**, 093002 (2011) [arXiv:1103.1366 [hep-ph]].

[20] J. Kopp, P. A. N. Machado and S. J. Parke, Phys. Rev. D **82**, 113002 (2010) [arXiv:1009.0014 [hep-ph]].

[21] N. C. Ribeiro, H. Nunokawa, T. Kajita, S. Nakayama, P. Ko and H. Minakata, Phys. Rev. D **77**, 073007 (2008) [arXiv:0712.4314 [hep-ph]].

[22] P. Huber, M. Lindner, M. Rolinec and W. Winter, Phys. Rev. D **74**, 073003 (2006) [arXiv:hep-ph/0606119].

[23] J. Barranco, O. G. Miranda and T. I. Rashba, JHEP **0512**, 021 (2005) [arXiv:hep-ph/0508299].

[24] B. Kayser, E. Fischbach, S. P. Rosen and H. Spivack, Phys. Rev. D **20**, 87 (1979).

[25] R. C. Allen *et al.*, Phys. Rev. D **47**, 11 (1993).

[26] A. B. Balantekin and K. O. Ozansoy, Phys. Rev. D **76**, 095014 (2007) [arXiv:0710.0028 [hep-ph]].

[27] J. Barranco, A. Bolanos, O. G. Miranda, C. A. Moura and T. I. Rashba, Phys. Rev. D **79**, 073011 (2009) [arXiv:0901.2099 [hep-ph]].

[28] K. Cheung, W. Y. Keung and T. C. Yuan, Phys. Rev. D **76**, 055003 (2007) [arXiv:0706.3155 [hep-ph]].

[29] T. i. Hur, P. Ko and X. H. Wu, Phys. Rev. D **76**, 096008 (2007) [arXiv:0709.0629 [hep-ph]].

[30] H. M. Chang *et al.* [TEXONO Collaboration], Phys. Rev. D **75**, 052004 (2007) [arXiv:hep-ex/0609001].

[31] Th. A. Mueller, D. Lhuillier, M. Fallot, A. Letourneau, S. Cormon, M. Fechner, L. Giot, T. Lasserre, J. Martino, G. Mention, A. Porta, F. Yermia, arXiv:1101.2663v3 [hep-ex]

[32] M. C. Gonzalez-Garcia, P. C. de Holanda and R. Zukanovich Funchal, JCAP **0806**, 019 (2008) [arXiv:0803.1180 [hep-ph]].

[33] D. A. Dicus and E. W. Kolb, Phys. Rev. D **15**, 977 (1977).

[34] P. Sutherland, J. N. Ng, E. Flowers, M. Ruderman and C. Inman, Phys. Rev. D **13**, 2700 (1976).

[35] H. T. Wong, H. B. Li, J. Li, Q. Yue and Z. Y. Zhou, J. Phys. Conf. Ser. **39**, 266 (2006) [arXiv:hep-ex/0511001].

[36] J. I. Collar [CoGeNT Collaboration], J. Phys. Conf. Ser. **136**, 022009 (2008).

[37] A. J. Anderson, J. M. Conrad, E. Figueroa-Feliciano, K. Scholberg and J. Spitz, arXiv:1103.4894 [hep-ph].

[38] K. Scholberg, Phys. Rev. D **73**, 033005 (2006) [arXiv:hep-ex/0511042].